

Received: 2015.04.27
Accepted: 2015.08.19
Published: 2015.12.31

MSX1 gene in the etiology orofacial deformities

Gen MSX1 w etiologii wad rozwojowych twarzoczaszki

Anna Paradowska-Stolarz

Department of Dentofacial Anomalies, Department of Orthodontics and Dentofacial Orthopedics, Wrocław Medical University, Poland

Summary

The muscle segment homeobox (*MSX1*) gene plays a crucial role in epithelial-mesenchymal tissue interactions in craniofacial development. It plays a regulative role in cellular proliferation, differentiation and cell death. The human *MSX1* domain was also found in cow (*Bt 302906*), mouse (*Mm 123311*), rat (*Rn13592001*), chicken (*Gg 170873*) and clawed toad (*XI 547690*).

Cleft lip and palate is the most common anomaly of the facial part of the skull. The etiology is not fully understood, but it is believed that the key role is played by the genetic factor activated by environmental factors. Among the candidate genes whose mutations could lead to formation of the cleft, the *MSX1* homeobox gene is mentioned. Mutations in the gene *MSX1* can lead to isolated cleft deformities, but also cause other dismorphic changes. Among the most frequently mentioned is loss of permanent tooth buds (mostly of less than 4 teeth – hypodontia, including second premolars).

Mutations of *MSX1* are observed in the Pierre- Robin sequence, which may be one of the features of congenital defects or is observed as an isolated defect. Mutation of the gene can lead to the occurrence of a rare congenital defect Wiktop (dental-nail) syndrome. Deletion of a fragment *MSX1* (4p16.3) located in the WHS critical region, may be a cause of some symptoms of Wolf-Hirschhorn syndrome.

Key words: **MSX1 homeobox • cleft lip and/or palate • hypodontia**

Full-text PDF: <http://www.phmd.pl/fulltxt.php?ICID=1192170>

Word count: 2144
Tables: –
Figures: –
References: 65

Author's address: Anna Paradowska-Stolarz, Ph.D., Department of Dentofacial Anomalies, Department of Orthodontics and Dentofacial Orthopedics, Wrocław Medical University, Krakowska 26, 50-425 Wrocław; e-mail: anna.paradowska-stolarz@umed.wroc.pl

The *MSX1* (muscle segment homeobox) gene plays a crucial role in epithelial-mesenchymal tissue interactions in craniofacial development. It plays a regulative role in cellular proliferation, differentiation and cell death – the processes playing a key role in cell growth and morphogenesis [4]. The *MSX1* gene, as well as *PAX9*, *AXIN2*, *EDA* and *EDAR* genes is responsible for tooth hypodontia [35]. The *MSX1* gene is located on chromosome 4p16.1. Its mutation causes autosomally dominant agenesis of second premolars and third molars. The gene causes multiple tooth agenesis, that is a result of a change in nucleotide sequence, where proline substitutes arginine in position 196 [24]. According to Modesto et al. [37], *MSX1* mutation leads to cleft lip and palate formation, when it occurs in parallel with the *PAX9* gene mutation. The mutations within *MSX1* may also lead to rare congenital defects such as Wiktop syndrome and Wolf-Hirschhorn syndrome. It has previously been shown that *MSX1* mutations notably change the quality of dental mesenchyme, but do not have its expression in dental epithelia at the bud, cap or bell stages of dental development [60].

The human *MSX1* domain was also found in cow (*Bt* 302906), mouse (*Mm* 123311), rat (*Rn*13592001), chicken (*Gg* 170873) and clawed toad (*XI* 547690) [23].

NONSyndromic Cleft Deformities

Cleft deformities are the most common facial malformation and are observed in ca. 10% of all the facial deformities [18]. Clefts of lip, alveolar bone and palate are observed two to three times more frequently in males while as isolated cleft of the palate is more frequent in females [54,56]. The formation of facial clefts takes place within 5-12 week of fetal development, during face formation [38]. The cleft lip and/or palate is a common congenital disorder, observed in 1:700 live births. The full etiology is not known, although a genetic factor is indisputable. There are numerous genes involved in palate formation, which disturbances of which may lead to cleft formation [44].

Observation of cleft's familial occurrence and selective presence in only some of the individuals indicates the genetic basis of the deformity, but also suggests coexistence of other factor. The etiology of clefts is multifactor, the genetics is only a basis "triggered" by environmental causes. The genetic factor is thought to play a crucial role in 20% of cases of clefts [1,43,54]. The genetic factors are divided as follows: genes encoding signaling molecules (*Bmp2*, *Bmp4*, *Bmp7*, *Shh*, *Wnt5a*, *Smad 2-4*), growth factors (*Egf*, *Egfr*, *Fgf1*, *Fgf2*, *Fgf8*, *Tgfa*, *Tgfβ1-3*) and their receptors (*Fgfr1*, *Fgfr2*), transcription factors (*Pax9*, *Tbx22*, *Msx1*, *Tbx1*, *Ap2α*, *Blx1-6*, *Lhx6*, *Gli 2-3*, *Hoxa2*, *Irf6*, *Pitx1*, *Pitx2*, *Prx1*), cellular adhesion molecules (*Pvrl1*, cadherin E, connexin 43), and extracellular matrix (fibronexin, *Col 1A2*, *Col2A1*, *Col11A1*, *Mmp2*, *Mmp3*, *Mmp9*, *Mmp13*, *Timp1-3*) [46,49]. Beside the multigenetic background, the *MSX1* and *TGFβ3* genes are found to be the genes most strongly related to cleft anomalies [59]. Many other genes related to the

clefts have their *loci* in different regions of chromosome 6, among them 6p23-24 and 6p24.3 [8].

Probably, most of the cases of isolated clefts are associated with either *MSX1* or *IRF6* mutation – it is assumed that the relation is observed in one third of cases [51]. The other candidate gene for cleft deformity might be *MTHFR* mutation of which lead to disturbances in folic acid transformation and storage [21]. *MSX1* is responsible mainly for secondary palate formation, therefore mutations within that gene is responsible for isolated cleft palate. *MSX1* is responsible for 2% of all non-syndromic clefts. Due to the complex cause of cleft deformities and to gene-to-gene interaction the influence on cleft palate formation is 9.7 times higher, which means it plays role in almost 20% of cases of clefts [29].

An association between *MSX1*-CA marker has been found. The clefts were more frequently observed in the *MSX1* homozygous allele 4 individuals, whose mothers smoked cigarettes or with both parents smoking. Similar results were obtained in children of mothers without daily folic supplementation during the periconceptual period [5].

The strongest association between nonsyndromic clefts and gene mutation is observed for the rs6446693 region in the *MSX1* gene [21]. Five variants of *MSX1* mutation were found as responsible for cleft palate forming. There were: A34G, G110G, P147Q, M37L and G267A. Among them, the rarest polymorphic variant is G267A mutation [55]. However, the thesis that *MSX1* is a factor influencing clefts is not promoted by all the reearchers. Some claim that *MSX1* might not be an etiologic factor responsible for orofacial clefting [12,30,36,37]. Therefore, further genetic analysis of other genes involved in facial development is required.

Wolf-Hirschhorn Syndrome

Wolf-Hirschhorn syndrome (WHS), also known as chromosome microdeletion 4p syndrome or Dillian syndrome, is a rare congenital disorder observed in 1:50,000 live births. It was first described in 1961 by two independent researchers: Hirschhorn and Cooper. The deformity is more frequently observed in women. Wolf-Hirschhorn syndrome is probably caused by mutation of one of two candidate genes (*WHS1* and *WHS2*). The syndrome is most frequently associated with a deletion within the 4p16.3 chromosome region (165kb), which encodes the *MSX1* gene [17,47,50,61,63,65]. The deletion most probably refers to *HOX7* gene, that maps to 4p16.1 gene [20]. The majority of cases is caused by "pure" *de novo* deletion and the greater the chromosomal material loss, the greater expression of Wolf-Hirschhorn syndrome [48,50].

Children suffer from multiple disorders, which include developmental delay and high mortality range of 30%. The patients suffer from congenital heart defects, muscle hypotonia, urinary tract malformations, cochlear hearing loss and immunodeficiency. The malformations



are so much severe that they may lead to patient's death [11,16,33,47].

The facial features include high forehead with prominent *glabella*, hypertelorism (with “protruding” eyes), high arched eyebrows, broad nasal bridge and short philtrum. A characteristic feature is micrognathia. In some cases they can be associated with facial asymmetry. The head and face take the characteristic look of a “Greek warrior helmet”. The mouth has short upper lip and downturned corners [3,41,47].

Multiple tooth agenesis occurring in this syndrome affects premolars and molars mainly. It is associated with *MSX1* mutation and may vary much (from multiple tooth agenesis to oligodontia). Other dental anomalies include taurodontism and late dental development [2,24,25,37,40].

PIERRE ROBIN SYNDROME

Pierre-Robin syndrome (or Pierre-Robin Sequence) is known from the triad of symptoms: congenital micrognathia, cleft of secondary palate and *glossoptosis* with upper airway obstruction. In some cases U-shaped palate and obstructive apnea are observed. The occurrence is relatively rare and ranges from 1:8,000 to 1:30,000 births. The sequence might be isolated or be associated with various congenital syndromes, eg. Treacher-Collins syndrome [13,42]. Beside the three characteristics, other defects might be observed. Those include dental anomalies, in particular hypodontia. This anomaly is observed most frequently in mandibular second premolars [31]. Hypodontia in this region accentuates mandibular undergrowth, as loss of tooth buds is a factor of bone underdevelopment in this area [53].

The primary anomaly of Pierre Robin syndrome is failure of mandibular growth caused by a failure in dorsal hold of the head in embryonic development. As a consequence, this forces high position of the tongue in primordia (primary oral cavity). The final result of that is physical obstruction for the posterior palatal shelves to fuse [42]. Micrognathia is found to be connected with deletions within genes 4p16-14, 4q31-35, 6q25-27 and 11q23. Those genes might be associated with Pierre Robin sequence, though its real etiology (and its probable genetic basis) is not known. The most probable genes involved in this syndrome are: the transcription factor *SOX9* (on 17q24.3-q25.1), *GAD67* (on 2q31) and *PVRL1* (on 11q23-q24) [22]. The *SOX9* mutation alters binding of the transcription factor *MSX1* [22,58]. In experiment performed in mice, it had been proved that lacking of *MSX1* function results in cleft of secondary palate, micrognathia and micrognathia, which are the most characteristic features of Pierre Robin syndrome as well. Furthermore, the mutation of this transcription factor causes underdevelopment of nasal, frontal and parietal bones, and the malleus. It also disturbs tooth development [45]. In families where *SOX9* mutation is the cause of Pierre Robin syndrome, the sequence is more severely expressed [42].

WIKTOP SYNDROME

Wiktop syndrome is a rare congenital disease, inherited autosomally dominantly. It is also known as tooth and nail syndrome (TNS) or “nail dysgenesis and hypodontia”, and as a type of ectodermal dysplasia. Tooth agenesis is accompanied by nail deformities - the nail plates are defective and thinner [26]. It was firstly described by Wiktop in 1965 [62] and its prevalence is estimated to ca. 1-2:10.000 births [22].

The syndrome is inherited autosomally dominantly [10]. The symptoms are less pronounced than in other types of ectodermal dysplasia, which is why those can be missed by the clinicians [64]. In 2001 it had been proved that the mesenchyme gene *MSX1* is responsible for Wiktop syndrome [26]. In 2013 the 3'-UTR region of this gene was found to be the most predominant locus [14].

In this congenital disorder, nails are dysplastic, thin, spoon-shaped (*koilonychia*) and easy to break (*onychorrhexis*). They grow slowly. Changes are more pronounced in toenails than fingernails. In some cases, nail defects are less pronounced in adulthood and less characteristic [15,19,52,55,64].

The other characteristic feature is hypodontia. There is no pattern of missing teeth - it is dependent on the individuals [19]. The most frequently missing teeth are lower incisors, secondary molars and upper canines [after 10]. The teeth (if present) show deformities such as peg-shape (conical shape). They can also be tapered or pointed [19]. The teeth are narrower and their eruption is disturbed [55].

Due to tooth agenesis, prosthetic appliances to improve chewing function and esthetic appearance might be required. In some cases they can also play a role as space maintainers [55]. Due to the hypodontia, the alveolar bone is hypoplastic and narrowed in its vertical dimension. The jaws are underdeveloped. Loss of occlusion in the vertical dimension may lead to lip eversion [19].

DENTAL ANOMALIES

Along with the *PAX9* gene, *MSX1* homeodomain is responsible for tooth agenesis. Both of them are expressed in dental mesenchyme. When homozygously deleted, they result in tooth development delay in an early stage. *MSX1* and *PAX9* mutations lead to disturbances of tooth development in the cap and/or bud stages of dental development. In mice, heterozygous deletion of either *PAX9* or *MSX1*, has no expression in dental structure. However, double heterozygous mutation shows arrested tooth development. This could be still rescued by Bmp4-signaling factor expression. Unlike in mice, in humans a heterozygous mutation in either *MSX1* or *PAX9* leads to tooth agenesis, predominantly in the posterior area. The mutation of *MSX1* leads to lack of second premolars and third mo-

lars, while *PAX9* mutation is responsible for lack of “wisdom teeth” [6,28,29,59]. Although *MSX1* mutation is the most frequent cause of lack of tooth buds, the mutation in A240P (rs4904210) in the *PAX9* gene results in multiple tooth agenesis, called oligodontia [39].

The nonsense mutation within the *MSX1* gene (eg. c.C565T in exon 2 or intron 1) dramatically reduces mRNA expression, suggesting that it may lead to rapid degradation of the mutated transcript, and cause the tooth agenesis [6,34]. Hypodontia (lack of 4 or fewer tooth buds) may also be a result of heterozygous *R151S* mutation. The allele is mildly deleterious for familial hypodontia. The mutation within this allele itself is insufficient to cause orofacial clefting. When other factors are involved (environmental or other, genetic ones), the cleft might occur [32]. It has also been shown that *MSX1* gene mutation (Ala219Thr, specifically), might be a reason for oligodontia, though due to the fact that lack of more than 4 tooth buds is more characteristic for other genes mutations, further studies of that topic are required [7,12,27]. There are studies, however, indicating that *MSX1* mutation is not a factor causing hypodontia. They point to *WNT10A* as a major cause of isolated congenital lack of teeth [57].

Beside tooth agenesis, underdevelopment of nails (its thinner and defective structure) in *MSX1*-null mice was observed [26]. Expression of the gene is unequal in the individuals and families. It takes place not only in mesenchyme structures, but also in the neural crest, cranial sensory placodes and bones [6,9]. Other developmental anomalies could be a result of *MSX1* mutation as well. Those could be deformities such as choanal atresia, anophthalmia, microphthalmia, cataracts, pachygyria, depigmentation and mental retardation were observed in mice with *MSX1* mutated gene. It might be the main responsibility of *MSX1* gene to modulate the growth and inhibit differentiation. The outgrowth could be reduced by the early differentiation in progress zone of facial processes [1].

Summing up, *MSX1* is an important gene expressing in mesenchyme. Its mutations may lead to developmental disorders such as variety of clefts, tooth and nail anomalies, and many others. The *MSX1* mutation is observed in Pierre-Robin sequence. The mutations may lead to Witkop syndrome or Wolf-Hirschhorn syndrome. Further research is required to establish the exact role of this gene.

REFERENCES

- [1] Abu-Hussein M.: Cleft lip and palate – etiological factors. *Dent. Med. Probl.*, 2012; 49: 149-156
- [2] Babich S.B., Banducci C., Teplitsky P.: Dental characteristics of the Wolf-Hirschhorn syndrome: a case report. *Spec. Care Dentist.*, 2004; 24: 229-231
- [3] Battaglia A., Filippi T., Carey J.C.: Update on the clinical features and natural history of Wolf-Hirschhorn (4p-) syndrome: experience with 87 patients and recommendations for routine health supervision. *Am. J. Med. Genet. C Semin. Med. Genet.*, 2008; 148C: 246-251
- [4] Bendall A.J., Abate-Shen C.: Roles for *Msx* and *Dlx* homeoproteins in vertebrate development. *Gene*, 2000; 247: 17-31
- [5] Benko S., Fantes J.A., Amiel J., Kleinjan D.J., Thomas S., Ramsay J., Jamshidi N., Essafi A., Heaney S., Gordon C.T., McBride D., Golzio C., Fisher M., Perry P., Abadie V., Ayuso C., et al.: Highly conserved non-coding elements on either side of *SOX9* associated with Pierre Robin sequence. *Nat. Genet.*, 2009; 41: 359-364
- [6] Brook A.H.: Multilevel complex interactions between genetic, epigenetic and environmental factors in the aetiology of anomalies of dental development. *Arch. Oral Biol.*, 2009; 54: S3-S17
- [7] Chishti M.S., Muhammad D., Haider M., Ahmad W.: A novel missense mutation *MSX1* underlines autosomal recessive oligodontia with associated dental anomalies in Pakistani families. *J. Hum. Genet.*, 2006; 51: 872-878
- [8] Cobourne M.T.: The complex genetics of cleft lip and palate. *Eur. J. Orthod.*, 2004; 26: 7-16
- [9] Davidson D.: The function and evolution of *Msx* genes: pointers and paradoxes. *Trends Genet.*, 1995; 11: 405-411
- [10] Devadas S., Varma B., Mungara J., Joseph T., Saraswathi T.R.: Witkop tooth and nail syndrome: a case report. *Int. J. Paediatr. Dent.*, 2005; 15: 364-369
- [11] Elder F.F., Colasurdo G.N., Rose V.M.: Wolf-Hirschhorn-Pitt-Rogers-Danks phenotype with no detectable deletion. *Am. J. Hum. Genet.*, 2000; 67; suppl. 2: 119
- [12] Etheredge A.J., Christensen K., Del Junco D., Murray J.C., Mitchell L.E.: Evaluation of two methods for assessing gene-environment interactions using data from the Danish case-control study of facial clefts. *Birth Defects Res. A Clin. Mol. Teratol.*, 2005; 73: 541-546
- [13] Figueroa A.A., Glupker T.J., Fitz M.G., BeGole E.A.: Mandible, tongue and airway in Pierre Robin sequence: a longitudinal cephalometric study. *Cleft Palate Craniofac. J.*, 1991; 28: 425-434
- [14] Ghaderi F., Hekmat S., Ghaderi R., Fardeh M.: *MSX1* mutation in Witkop syndrome; a case report. *Iran J. Med. Sci.*, 2013; 38: 191-194
- [15] Giansanti J.S., Long S.M., Rankin J.L.: The “tooth and nail” type of autosomal dominant ectodermal dysplasia. *Oral Surg. Oral Med. Oral Pathol.*, 1974; 37: 576-582
- [16] Hanley-Lopez J., Estabrooks L.L., Stiehm R.: Antibody deficiency in Wolf-Hirschhorn syndrome. *J. Pediatr.*, 1998; 133: 141-143
- [17] Hirschhorn K., Cooper H.L.: Apparent deletion of one chromosome (4 or 5) in a child with defects of midline fusion. *Hum. Chrom. News*, 1961; 4: 14
- [18] Hozyasz K.: Wpływ czynników środowiskowych na występowanie wad rozszczepowych twarzoczaszki. *Med. Rodz.*, 1999; 2: 11-17
- [19] Hudson C.D., Witkop C.J.: Autosomal dominant hypodontia with nail dysgenesis. Report of twenty-nine cases in six families. *Oral Surg. Oral Med. Oral Pathol.*, 1975; 39: 409-423
- [20] Ivens A., Flavin N., Williamson R., Dixon M., Bates G., Buckingham M., Robert B.: The human homeobox gene *HOX7* maps to chromosome 4p16.1 and may be implicated in Wolf-Hirschhorn syndrome. *Hum. Genet.*, 1990; 84: 473-476
- [21] Jagomägi T., Nikopensius T., Kjutskov K., Tammekivi V., Viltrop T., Saag M., Metspalu A.: *MTHFR* and *MSX1* contribute to the risk of nonsyndromic cleft lip/palate. *Eur. J. Oral Sci.*, 2010; 118: 213-220
- [22] Jakobsen L.P., Knudsen M.A., Lespinasse J., Ayuso C.G., Ramos C., Fryns J.P., Bugge M., Tommerup N.: The genetic basis of Pierre Robin Sequence. *Cleft Palate Craniofac. J.*, 2006; 43: 155-159



- [23] Jezewski P.A., Vieira A.R., Nishimura C., Ludwig B., Johnson M., O'Brien S.E., Daack-Hirsch S., Schultz R.E., Weber A., Nepomucena B., Romitti P.A., Christensen K., Orioli I.M., Castilla E.E., Machida J., Natsume N., Murray J.C.: Complete sequencing shows a role for MSX1 in non-syndromic cleft lip and palate. *J. Med. Genet.*, 2003; 40: 399-407
- [24] Jędrzysek A., Kmiecik M., Paszkiewicz A.: Review of modern knowledge on hypodontia. *Dent. Med. Probl.*, 2009; 46: 118-125
- [25] Johnston N.J., Franklin D.L.: Dental findings of a child with Wolf-Hirschhorn syndrome. *Int. J. Paediatr. Dent.*, 2006; 16: 139-142
- [26] Jumlongras D., Bei M., Stimson J.M., Wang W.F., DePalma S.R., Seidman C.E., Felbor U., Maas R., Seidman J.G., Olsen B.R.: A nonsense mutation in MSX1 causes Witkop syndrome. *Am. J. Hum. Genet.*, 2001; 69: 67-74
- [27] Kamamoto M., Machida J., Yamaguchi S., Kimura M., Ono T., Jezewski P.A., Higashi Y., Nakayama A., Shimozato K., Tokita Y.: Clinical and functional data implicate the Arg(151)Ser variant of MSX1 in familial hypodontia. *Eur. J. Hum. Genet.*, 2011; 19: 844-850
- [28] Kapadia H., Mues G., D'Souza R.: Genes affecting tooth morphogenesis. *Orthod. Craniofac. Res.*, 2007; 10: 105-113
- [29] Kohli S.S., Kohli V.S.: A comprehensive review of the genetic basis of cleft lip and palate. *J. Oral Maxillofac. Pathol.*, 2012; 16: 64-72
- [30] Koillinen H., Ollikainen V., Rautio J., Hukki J., Kere J.: Linkage and linkage disequilibrium searched for between non-syndromic cleft palate and four candidate loci. *J. Med. Genet.*, 2003; 40: 464-468
- [31] Larson M., Helidust R., Jakobsson O.P.: Dental abnormalities and ectopic eruption in patients with isolated cleft palate. *Scand. J. Plast. Reconstr. Surg. Hand Surg.*, 1997; 32: 203-212
- [32] Liang J., Zhu L., Meng L., Chen D., Bian Z.: Novel nonsense mutation in MSX1 causes tooth agenesis with cleft lip in a Chinese family. *Eur. J. Oral Sci.*, 2012; 120: 278-282
- [33] Lurie I.W., Lazjuk G.I., Ussova Y.I., Presman E.B., Burevich D.B.: The Wolf-Hirschhorn syndrome. *I. Genetics. Clin. Genet.*, 1980; 17: 375-384
- [34] MacKenzie A., Leeming G.L., Jowett A.K., Ferguson M.W., Sharpe P.T.: The homeobox gene Hox 7.1 has specific regional and temporal expression patterns during early murine craniofacial embryogenesis, especially tooth development in vivo and in vitro. *Development*, 1991; 111: 269-285
- [35] Mielnik-Błaszczak M., Piątkowska A., Machowska A., Bukowska B.: Tooth Genesis – a review of literature and own case studies. *Dent. Med. Probl.*, 2012; 49: 293-298
- [36] Mitchell L.E., Murray J.C., O'Brien S., Christensen K.: Evaluation of two putative susceptibility loci for oral clefts in the Danish population. *Am. J. Epidemiol.*, 2001; 153: 1007-1015
- [37] Modesto A., Moreno L.M., Krahn K., King S., Lidra A.C.: MSX1 and orofacial clefting with and without tooth agenesis. *J. Dent. Res.*, 2006; 85: 542-546
- [38] Moore G., Ivens A., Chambers J., Bjornsson A., Arnason A., Jenson O., Williamson R.: The application of molecular genetics to detection of craniofacial abnormality. *Development*, 1988; 103: 233-239
- [39] Mostowska A., Biedziak B., Jagodziński P.P.: Analysis of the coding sequence of the MSX1 and PAX9 genes in patients with congenital lack of permanent teeth. *Dent. Med. Probl.*, 2012; 49: 157-165
- [40] Nieminen P., Kotilainen J., Aalto Y., Knuutila S., Pirinen S., Theleff I.: MSX1 gene is deleted in Wolf-Hirschhorn syndrome patients with oligodontia. *J. Dent. Res.*, 2003; 82: 1013-1017
- [41] Pitt, D.B., Rogers, J.G., Danks, D.M.: Mental retardation, unusual face, and intrauterine growth retardation: a new recessive syndrome? *Am. J. Med. Genet.*, 1984; 19: 307-313
- [42] Rainger J.K., Bhatia S., Bengani H., Gautier P., Rainger J., Pearson M., Ansari M., Crow J., Mehendale F., Palinkasova B., Dixon M.J., Thompson P.J., Martin M., Sisodiya S.M., Kleinjan D.A., Fitzpatrick D.R.: Disruption of SATB2 or its long-range cis-regulation by SOX9 causes a syndromic form of Pierre Robin sequence. *Hum. Mol. Genet.*, 2014; 15: 2569-2579
- [43] Rychlik D., Wójcicki P., Koźlik M.: Osteoplasty of the alveolar cleft defect. *Adv. Clin. Exp. Med.*, 2012; 21: 255-262
- [44] Salahshourifar I., Halim A.S., Wan Sulaiman W.A., Zilfalil B.A.: Contribution of MSX1 variants to the risk of non-syndromic cleft lip and palate in a Malay population. *J. Hum. Genet.*, 2011; 56: 755-758
- [45] Satokata I., Maas R.: Msx1 deficient mice exhibit cleft palate and abnormalities of craniofacial and tooth development. *Nat. Genet.*, 1994; 6: 348-356
- [46] Schutte B.C., Murray J.C.: The many faces and factors of orofacial clefts. *Hum. Mol. Genet.*, 1999; 8: 1853-1859
- [47] Shannon N.L., Maltby E.L., Rigby A.S., Quarrell O.W.: An epidemiological study of Wolf-Hirschhorn syndrome: life expectancy and cause of mortality. *J. Med. Genet.*, 2001; 38: 674-679
- [48] Sifakis S., Manolagos E., Vetro A., Kappou D., Peitsidis P., Kontodiu M., Garas A., Vrachnis N., Konstantinidou A., Zuffardi O., Orru S., Papoulidis I.: Prenatal diagnosis of Wolf-Hirschhorn syndrome confirmed by comparative genomic hybridization array: report of two cases and review of the literature. *Mol. Cytogenet.*, 2012; 5: 12
- [49] Stanier P., Moore G.E.: Genetics of cleft lip and palate: syndromic genes contribute to the incidence of non-syndromic clefts. *Hum. Mol. Genet.*, 2004; 13: R73-R81
- [50] Stec I., Wright T.J., van Ommen G.J., de Boer P.A., van Haeringen A., Moorman A.F., Altherr M.R., den Dunnen J.T.: WHSC1, a 90 kb SET domain containing gene, expressed in early development and homologous to a Drosophila dysmorphia gene maps in the Wolf-Hirschhorn syndrome critical region and is fused to IgH in t(4;14) multiple myeloma. *Hum. Mol. Genet.*, 1998; 7: 1071-1082
- [51] Steponavičiūtė D., Kasnauskienė J., Kučinskas V.: Nucleotide sequence changes in the MSX1 and IRF6 genes in Lithuanian patients with nonsyndromic orofacial clefting. *Acta Med. Lithuanica*, 2006; 13: 216-225
- [52] Subramaniam P., Neeraja G.: Witkop's tooth and nail syndrome: a multifaceted approach to dental management. *J. Indian. Soc. Pedod. Prev. Dent.*, 2008; 26: 22-25
- [53] Suri S., Ross R.B., Tompson B.D.: Mandibular morphology and growth with and without hypodontia in subjects with Pierre Robin sequence. *Am. J. Orthod. Dentofac. Orthop.*, 2006; 130: 37-46
- [54] Szeląg J., Noga L., Orłowska K., Pałka Ł., Paradowska A.: Analiza wpływu endo- i egzogennych czynników ryzyka w etiologii rozszczepów podniebienia pierwotnego i wtórnego. *Dent. Med. Probl.*, 2006; 43: 556-562
- [55] Świder K., Szozda A., Tokarski T.: Prosthodontic treatment of children – case reports. *Dent. Med. Probl.*, 2013; 50: 106-113
- [56] Tortora C., Meazzini M.C., Garratini G., Brusati R.: Prevalence of abnormalities in dental structure, position and eruption pattern in a population of unilateral and bilateral cleft lip and palate patients. *Cleft Palate Craniofac. J.*, 2008; 45: 154-162
- [57] van den Boogaard M.J., Creton M., Bronkhorst Y., van der Hout A., Hennekam E., Lindhout D., Cune M., Ploos van Amstel H.K.: WNT10A, rather than MSX1 is a major cause of non-syndromic hypodontia. *J. Med. Genet.*, 2012; 49: 327-331
- [58] van den Boogaard M.J., de Costa D., Krapels I.P., Liu F., van Duin C., Sinke R.J., Lindhout D., Steegers-Theunissen R.P.: The MSX1 allele 4 homozygous child exposed to smoking at periconception is most sensitive in developing nonsyndromic orofacial clefts. *Hum. Genet.*, 2008; 124: 525-534

- [59] Vieira A.R., Orioli I.M., Castilla E.E., Cooper M.E., Marazita M.L., Murray J.C.: MSX1 and TGF β 3 contribute to clefting in south America. *J. Dent. Res.*, 2003; 82: 289-292
- [60] Wang J., Jjan F., Chen J., Wang H., Lin Y., Yang Z., Pan X., Lai W.: Sequence analysis of PAX9, MSX1 and AXIN2 genes in a Chinese oligodontia family. *Arch. Oral Biol.*, 2011; 56: 1027-1034
- [61] Wiczorek D., Krause M., Majewski F., Albrecht B., Horn D., Riess O., Gillessen-Kaesbach G.: Effect of the size of the deletion and clinical manifestation in Wolf-Hirschhorn syndrome: analysis of 13 patients with a de novo deletion. *Eur. J. Hum. Genet.*, 2000; 8: 519-526
- [62] Witkop C.J.: Genetic disease of the oralcavity. In: Tietze R.W. (ed.) *Oralpathology*. Blakiston Division, McGraw-Hill, New York, 1965, 810-814
- [63] Wright T.J., Rieke D.O., Denison K., Abmayr S., Cotter P.D., Hirschhorn K., Keinanen M., McDonald-McGinn D., Somer M., Spinner N., Yang-Feng T., Zackai E., Altherr M.R.: A transcript map of the newly defined 165kb Wolf-Hirschhorn syndrome critical region. *Hum. Mol. Genet.*, 1997; 6: 317-324
- [64] Zabawski E.J.Jr., Cohen J.B.: Hereditary hypodontia and onychorrhexis of the fingernails and toenail koilonychia: Witkop's tooth and nail syndrome. *Dermatol. Online J.*, 1999; 5: 3
- [65] Zollino M., Lecce R., Fischetto R., Murdolo M., Faravelli F., Selicorni A., Buttè C., Memo L., Capovilla G., Neri G.: Mapping the Wolf-Hirschhorn syndrome phenotype outside the currently accepted WHS critical region and defining a new critical region, WHSCR-2. *Am. J. Hum. Genet.*, 2003; 75: 590-597

The author has no potential conflicts of interest to declare.

